

Tabla 5: RN tratados con hipotermia

| | No Hipotermia | | | | Hipotermia | | | |
|-------|---------------|--------------------------|--------------------------|-------------------------|--------------|--------------------------|--------------------------|-------------------------|
| | P Pre 63,17 | | | | P Pre 47,88 | | | |
| | Estudios (N) | P post positiva (IC 95%) | P post negativa (IC 95%) | OR diagnóstica (IC 95%) | Estudios (N) | P post positiva (IC 95%) | P post negativa (IC 95%) | OR diagnóstica (IC 95%) |
| 1-30d | 13 (543) | 86% (80 a 90%) | 23% (25 a 30%) | 29,80 (17,09 a 51,95) | 4 (181) | 80% (74 a 86%) | 14% (8 a 24%) | 29,59 (12,12 a 72,25) |
| 1-7d | 6 (211) | 88 (76 a 95%) | 22 (9 a 42%) | 31,05 (10,62 a 90,84) | 2 (32) | 86% (62 a 96%) | 12% (1 a 78%) | 48,34 (1,85 a 1246,90) |
| 8-30d | 5 (149) | 88% (68 a 96%) | 20% (5 a 56%) | 43,87 (6,94 a 281,31) | 1 (36) | 73,5 (43,3 a 90,3%) | 28,3% (14,3 a 48,3%) | |