

Anexo 4. Abreviaturas

| | |
|---------|---|
| AAA | Ácido acetilsalicílico |
| ACEi | <i>Angiotensin-converting enzyme (ACE) inhibitors</i> |
| ACV | Accidente Cerebrovascular |
| AIT | Accidente Isquémico Transitorio |
| AP | Atención Primaria |
| ARA-2 | Antagonistas de los receptores de la angiotensina II |
| ARB | <i>Angiotensin receptor blockers</i> |
| AVAC | Año de vida ajustado por calidad |
| AVADs | Años de vida ajustados por discapacidad. |
| AVD | Actividades de la vida diaria |
| BPC | Buena Práctica Clínica |
| CAST | <i>Chinese Acute Stroke Trial</i> |
| CCBS | <i>Calcium channel blockers</i> |
| CIBIC | <i>Clinician Interview Based Impression of Change</i> |
| CNS-LS | Escala de labilidad del centro de estudios neurológicos (<i>Center for Neurologic Study-Lability Scale</i>) |
| CPSS | <i>Cincinnati Prehospital Stroke Scale</i> |
| CTAR | <i>Chin Tuck Against Resistance Exercise</i> |
| DDD | Dosis Diaria Definida |
| DE | Desviación Estándar |
| DEM | Diferencia Estandarizada de Medias |
| DM | Diferencia de Medias |
| E | Especificidad |
| EAN | <i>European Academy of Neurology</i> |
| EAT-10 | <i>Eating Assessment Tool-10</i> |
| ECA | Ensayo Clínico Aleatorizado |
| ECV | Enfermedad cardiovascular |
| EE | Error Estándar |
| ESO | <i>European Stroke Organization</i> |
| EMST | Entrenamiento de la fuerza muscular respiratoria (<i>Expiratory Muscle Strength Training</i>) |
| EQ | <i>EuroQol</i> |
| ERm | Estado Funcional |
| ESSD | <i>European Society for Swallowing Disorders.</i> |
| EtD | <i>Evidence to Decision</i> |
| EXPRESS | <i>Effect of urgent treatment of transient ischaemic attack and minor stroke on early recurrent stroke.</i> |

| | |
|----------------|--|
| <i>FAI</i> | <i>Frenchay Activity Index</i> |
| FAST | <i>Face, arms, speech, time test</i> |
| GLIAS | <i>GLyceria in Acute Stroke study</i> |
| GPC | Guía de Práctica Clínica |
| GRADE | <i>Grading of Recommendations Assessment, Development and Evaluation</i> |
| <i>GSES</i> | <i>General Self Efficacy Scale</i> |
| <i>HAMA</i> | <i>Hamilton Anxiety Rating Scale</i> |
| <i>HCISM</i> | <i>Hospital Community-integrated Service Model</i> |
| <i>HR</i> | <i>Hazard Ratio</i> |
| IC | Intervalo de Confianza |
| <i>ICaPPS©</i> | <i>Integrated care pathway for post-stroke patients</i> |
| IPA | <i>Impact on Participation and Autonomy</i> |
| <i>IST</i> | <i>International Stroke Trial</i> |
| LAPSS | <i>Los Angeles Prehospital Stroke Screening</i> |
| MASS | <i>Melbourne Ambulance Stroke Screen</i> |
| <i>MDPS</i> | <i>Medical Priority Dispatch System</i> |
| <i>MYR</i> | Ringgit malayo |
| NO | Óxido Nitrico (<i>Nitric oxid</i>) |
| <i>OR</i> | <i>Odds Ratio</i> |
| PAD | Presión Arterial Diastólica |
| PAS | Presión Arterial Sistólica |
| <i>PLSC</i> | <i>Pathological Laughter and Crying Scale</i> |
| QASC | <i>Quality in Acute Stroke Care</i> |
| RM | Resonancia Magnética |
| ROSIER | <i>Recognition of Stroke in the Emergency Room scale</i> |
| RR | Riesgo Relativo |
| RS | Revisión sistemática |
| S | Sensibilidad |
| sICH | Hemorragia Intracraneal Sintomática (<i>Symptomatic Intracranial Hemorrhage</i>) |
| SNS | Sistema Nacional de Salud |
| SSRI | <i>Selective serotonin reuptake inhibitors</i> |
| TC | Tomografía computerizada |
| TCA | Antidepresivos tricíclicos (<i>Tricyclic Antidepressives</i>). |
| TIV | Trombólisis Intravenosa |
| <i>TPRT</i> | <i>Tongue to Palate Resistance Training</i> |
| VPP | Valor Predictivo Positivo |